

# Population Doses from Gamma Radiation Exposure around Damaturu Metropolis, Yobe State, Nigeria

Habu Tela Abba\* and Agada Livinus Emeka

Received 2 May 2020/Accepted 25 May 2020/Published online: 26 May 2020

**Abstract** Humans are exposed to radiation in their environment with or without their consent; and the exposure to natural background radiation is an unpreventable event on earth. Exposure to terrestrial background gamma radiation in Damaturu metropolis was measured in 65 locations using a hand held radiation survey meter (RDS-31). The gamma absorbed dose rate in air 1 m above ground surface ranged from 5 to 165 nGy h<sup>-1</sup> with a mean value of 56±5 nGy h<sup>-1</sup>. The study results revealed that, higher dose rates are located around the main roads within the city. The mean value was found to be within the set limit by UNSCEAR for normal background radiation. Thus, the dose rates are classified as low radiation level and are below the level to cause acute health effects in humans. The results obtained would serve as baseline data for further studies on background radiation in the study area.

**Key Words:** Gamma radiation dose rate, isodose map, kriging method

\*Habu Tela Abba

Department of Physics, Yobe State University, Damaturu, Nigeria

Email: [htelaabba@gmail.com](mailto:htelaabba@gmail.com)

**Orcid id:** [0000-0003-0842-2235](https://orcid.org/0000-0003-0842-2235)

Agada Livinus Emeka

Department of Physics, Yobe State University, Damaturu, Nigeria

Email: [agadaman1908@gmail.com](mailto:agadaman1908@gmail.com)

**Orcid id:** [0000-0003-2884-8831](https://orcid.org/0000-0003-2884-8831)

## 1.0 Introduction

The global population is continually exposed to background radiation from both natural and anthropogenic sources. The natural radiation comes from cosmic and terrestrial radionuclides within the earth's crust mainly due to primordial radionuclides <sup>238</sup>U, <sup>232</sup>Th and <sup>40</sup>K that were created during the earth's formation (Al-Jundi, 2002). According to UNSCEAR (2000), 86% of the total radiation exposed to the global population are consequences of natural radionuclides while 14% is from anthropogenic sources. The dosage of exposure to gamma radiation from natural sources

is significantly determined by the amount of radionuclides in rocks and soil and vary from one location to another depending on the geology and geographical conditions of such locations (UNSCEAR, 2000). Therefore, exposure level to public varies from one geographical location to the other. Available literature revealed that people living in areas of granitic rock formation are more likely exposed to gamma radiation than those living in areas of sedimentary rocks indicating that different rock formation emits radiation at different rate (El-Arabi *et al.*, 2006). The worldwide annual effective dose received by the population from all natural and artificial sources is 2.8 mSv, out of which 85% of the dose (2.4 mSv) comes from only natural background radiations (Singh *et al.*, 2017).

Previous studies have shown that areas with high background radiation are found in Yangjiang, China; Kerele, India; and Ramsar, Iran (Ghiassi-Nejad *et al.*, 2002) and in Asia, maximum outdoor measurement was recorded in Malaysia and the maximum indoor measurement was recorded in Hong Kong and Iran (Gholami *et al.*, 2011). A study by Karahan and Bayulken (2000) reported mean exposure of 65 nGy<sup>-1</sup> for Istanbul with equivalent annual effective dose of 80 μS. In a similar study, Leung *et al.* (1990) reported that before the Chernobyl accident, gamma dose rate for Hong Kong was 0.076 nGy y<sup>-1</sup>. In a nationwide study, Furukawa and Shingaki (2012), found the average dose in Japan to be about 50 nGy y<sup>-1</sup>. In Nigeria, several studies have been conducted to measure gamma radiation level in some areas. For instance, it is reported that the mean outdoor gamma radiation dose rate was 250 nGy y<sup>-1</sup> in Jos, Plateau state (Abba *et al.*, 2017). Farai and Vincent (2006), reported that the equivalent dose due to outdoor exposure to radiation in Abeokuta, Nigeria ranged from 0.19 to 1.64 mSv/y with mean concentration of 0.45 mSv y<sup>-1</sup>. A nationwide study of the terrestrial radiation in Nigeria (Farai and Jibiri, 2000) indicates that the mean annual effective dose equivalent is 0.27 mSv y<sup>-1</sup>. Growth in population and technological advancement has led to the use and improper disposal of radioactive materials in

the environment. These activities will eventually enhance background concentrations of radiation and hence the amount and intensity that is emitted to the environment. Thus, studies of environmental natural gamma radiation level are significant in establishing baseline information on the geology and background level of radiation in such location. It helps in keeping data for normal background radiation before any eventualities in terms of radioactive contamination. Radiation data are useful tracers for atmospheric variational studies of environmental radioactivity and also furnish information that are relevant for evaluating national and international average values of radiometric and dosimetric quantities. It can also be used to check anthropogenic contamination of the environment (Ramli *et al.*, 2014).

Most of the studies conducted in the country are in the southern part of Nigeria with none or little in the northern region. Therefore, the present work aims at extending the survey to the towns in the northeastern region of Nigeria; it is also aimed at producing isodose map for the spatial distribution of background gamma radiation dose rates for the

metropolis. The result of this survey would serve as baseline data for future references. The result is also required for producing radiological map for the region.

## 2.0 Material and methods

### 2.1 Study area

Damaturu is the state capital of Yobe state and is located between latitudes  $11^{\circ}39'30'' - 11^{\circ}47'00''$  N and longitudes  $11^{\circ}34'00'' - 12^{\circ}02'00''$  E (Fig.1). The town is on A3 highway and covers a land area of 2,366 km<sup>2</sup>. According to 2006 census, the population of the area was projected at 69,952 people for 2010 (NPC, 2006). Geologically, Damaturu is located within the fringe of Chad basin, The Chad basin constitute of sedimentary rocks such as clay, sands, sand stones, and shale intercalations (Agada *et al.*, 2020). The climate of Damaturu is characterised by short wet season (June – September) and long dry season (October – May), with high temperatures of about 39°C to 49°C. During the raining season, temperatures fall to about 25°C with annual rainfall of about 500 to 1000 mm.





Fig. 1: Map of Yobe State showing study area where assessments were conducted.



### 2.2 Measurement of gamma radiation dose rate

A hand held modular radiation survey meter (RDS-31), was used to measure outdoor gamma radiation dose rates randomly at 65 different locations across the entire study area (Fig. 2). The instrument is versatile radiation detector and survey meter designed for a wide range of applications. Readings were displayed in  $\mu\text{S h}^{-1}$  which were subsequently converted to gamma

absorbed dose per hour ( $1 \text{ nGy h}^{-1} = 0.001 \mu\text{S h}^{-1}$ ). Measurements were taking in open fields away from buildings and other structure to minimize interference from artificial sources. In order to minimize systematic error, the readings were taken in triplicate at each location and the average was recorded. Measurements were conducted at 1 m above the ground surface and the coordinates of the sampling locations were recorded using a Global Positioning System (GPS) device.



**Fig. 2: Locations of gamma dose rates measurement**

### 3.0 Results and Discussion

Table 1 shows the descriptive statistics of gamma radiation dose rates readings. The results indicate that dose rate in Damaturu metropolis ranged from  $5 \text{ nGy h}^{-1}$  to  $165 \text{ nGy h}^{-1}$  with a mean value of  $56 \pm 5 \text{ nGy h}^{-1}$ . The lowest dose rate was recorded in the northeastern part of the town, an area with low population density and less activities while the highest measured dose rate was measured in central part of the town, an area that is densely populated with significant human activities including heavy movement of automobiles.

Enhanced gamma radiation levels were observed around the major roads within the metropolis. The results obtained indicate that background gamma radiation in Damaturu is below the UNSCEAR (2000) limit of  $59 \text{ nGy h}^{-1}$  for normal background gamma radiation. It also indicates that the natural

background radiation level in the city does not pose any significant health risk to the inhabitants. Nevertheless, sources of higher radiation, especially around the major roads should be checked in order to minimize future induced risk



through additional emission. The results agreed with previous studies conducted in some cities in Nigeria (Agba *et al.*, 2006; Farai and Vincent, 2006; Obioha and Okonkwo, 2001; Ramli, *et al.*, 2014).


**3.1 Isodose mapping of gamma radiation dose rates**

Kriging method of interpolation using geographical information system (GIS) software was applied to plot the projection of dose rates for the entire study area. The Kriging provides the best unbiased linear estimation of dose-rate values (Sanusi *et al.*, 2014). Fig. 3 provides the spatial distribution of gamma radiation dose rates in air 1 m above the ground in Damaturu metropolis. It was observed that higher levels of dose rates are concentrated at the centre of the town and two locations in the northern part of the study area. This indicates the possibility of an anthropogenic

sources in those areas with enhanced background radiation.

**Table 1: Descriptive statistics of the measured gamma dose rates**

Mean	56
Standard Error	5
Median	40
Mode	56
Standard Deviation	44
Kurtosis	0
Skewness	1
Range	160
Minimum	5
Maximum	165
Confidence Level (95%)	11



**Fig. 3: Isodose Map of Damaturu Metropolis**

**4.0 Conclusion**

This study present baseline information on the background gamma radiation dose rates in Damaturu and reveals that the lowest radiation level is in the northeastern zone of the city while the highest radiation value was observed in an area around the central market. Locations around

the major roads were observed to exhibit higher dose rates. The results of this study are within the regulatory limits and are within the range of the average readings of radiation limits reported in literature. Results from the study area indicated that the natural background radiation level in the city does not pose any significant health risk to the



inhabitants. The results provide the essential baseline information for the assessment of any environmental radioactive contamination of the area in foreseeable future. It is therefore recommended that further studies should be carried out to identify the source of high dose rates within the city centre and around the major roads.

### 5.0 References

- Abba, H., Saleh, M., Hassan, W., Aliyu, A. and Ramli, A. (2017). Mapping of natural gamma radiation (NGR) dose rate distribution in tin mining areas of Jos Plateau, Nigeria. *Environmental Earth Sciences*. 76(5), (208) PP. 201-209
- Agba, E. H., Onjefu, S. A. and Ugwuani, J. U. (2006). Preliminary investigation of ambient radiation levels of the of mining sites in Benue State, Nigeria. *Nig Journ Phys*, 18 (3), PP. 219-222. .
- Al-Jundi, J. (2002). Population doses from terrestrial gamma exposure in areas near to old phosphate mine, Russaifa, Jordan. *Radiation Measurements*. 35(1), PP. 23-28.
- El-Arabi, A., Abbady, A. G. and Hussein, A. (2006). Gamma-ray measurements of natural radioactivity in sedimentary rocks from Egypt. *Nuclear Science and Techniques*. 17(2), PP. 123-128.
- Farai, I. and Jibiri, N. (2000). Baseline studies of terrestrial outdoor gamma dose rate levels in Nigeria. *Radiation Protection Dosimetry*. 88(3), PP. 247-254.
- Farai, I. and Vincent, U. (2006). Out-door radiation level measurement in Abeokuta, Nigeria, by thermoluminescent dosimetry. *Nigerian Journal of Physics*. 18(1), PP. 121-126.
- Furukawa, M. and Shingaki, R. (2012). Terrestrial gamma radiation dose rate in Japan estimated before the 2011 Great East Japan Earthquake. *Rad Emergency Med*. 1(6), PP.65-72.
- Nejad, M., Mortazavi, S., Cameron, J., Niroomand-Rad, A. and Karam, P. (2002). Very high background radiation areas of Ramsar, Iran: preliminary biological studies. *Health physics*. 82(1), PP. 87-93.
- Gholami, M., Mirzaei, S. and Jomehzadeh, A. (2011). Gamma background radiation measurement in Lorestan province, Iran. 9(2), PP. 89-03.
- Karahan, G. and Bayulken, A. (2000). Assessment of gamma dose rates around Istanbul (Turkey). *Journal of Environmental Radioactivity*. 47(2), PP. 213-221.
- Leung, K., Lau, S. and Poon, C. (1990). Gamma radiation dose from radionuclides in Hong Kong soil. *Journal of environmental radioactivity*. 11(3), PP. 279-290.
- NPC (2006). National population Commission (NPC):. *Provisional of 2006 Census Results*. Abuja, Nigeria.
- Obioha, F. I. and Okonkwo, P. O. (2001). Background gamma radiation levels in the Nigerian environment. *West African Journal of Radiology*. 8(1), PP. 16-19.
- Ramli, A. T., Aliyu, A., Agba, E. and Saleh, M. (2014). Effective dose from natural background radiation in Keffi and Akwanga towns, Central Nigeria. *International Journal of Radiation Research*. 12(1), PP. 47-52.
- Sanusi, M., Ramli, A., Gabdo, H., Garba, N., Heryanshah, A., Wagiran, H. and Said, M. (2014). Isodose mapping of terrestrial gamma radiation dose rate of Selangor state, Kuala Lumpur and Putrajaya, Malaysia. *Journal of environmental radioactivity*. 135, PP. 67-74.
- UNSCEAR (2000). Sources and effects of ionizing radiation: . UNSCEAR 1993 report to the General Assembly with scientific annexes / United Nations Scientific Committee on the Effects of Atomic Radiation.: New York: United Nations.

