

Chemical Information from GCSM Analysis of Acetone Extract of *Piper guineense* Leaves. Part 1

Onyeije Ugomma Chibuzo* and Imeh Okop

Received 25 June 2020/Accepted 29 July 2020/Published online: 30 July 2020

Abstract Applications of plant leaves for various purposes is based on its chemical constituents which may include proximate, elemental, phytochemical, toxicant, amino acid and other toxicants. Knowledge of phytochemical constituents is significant for their pharmaceutical/medicinal values. This study was carried out to investigate the chemical constituents of acetone extract of *Piper guineense* leaves through phytochemical screening and GCMS analysis. Results obtained from phytochemical screening indicated the major constituents (those whose concentrations were greater than 1%) to include 6-dimethyloxazolo(5,4-c)pyridazin-4-amine (31.80 %), 3-(1-methylethyl)-cyclohexene (20.99 %), 4-methoxy-N-(4-nitrobenzyl)-benzamide (12.82 %), alpha bisabolene (7.33%), 1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methylene-1-(1-methylethyl)-naphthalene (4.42 %), 3,7-diacetamidophenoxathin (4.10 %), 1,3,3-trimethyl tricyclo{2,2,1}heptane (2.98%), 3H-indazol-3-one (2.11%), 1H-indene, octahydro-1,7a-dimethyl-4-(1-methylethenyl)-1,4-methano-1H-indene (1.98%), piperidine (1.97%), 2,4-disopropenyl-1-methyl-1-vinyl (1.70%), n-hexadecanoic acid (1.68%), eudesma-4[14],11-diene (1.27%). The pharmaceutical values of the identified constituents were also analysed. The study reveals that acetone extract of *Piper guineense* contains constituents that are not visible with some other solvents

Key Words: *Piper guineense* leaves, Chemical constituents, phytochemicals, GCMS, phytochemical screening

Onyeije Ugomma Chibuzo

Department of Chemistry

Rhema University, 153-155 Aba Owerri Road

P. M. B. 7021, Abia State, Nigeria

Email: ugommaoyeije@yahoo.com

Orcid id: [0000-0003-2832-7948](https://orcid.org/0000-0003-2832-7948)

Imeh Okop

Department of Chemistry

Akwa Ibom State University

Ikot Akpaden, Akwa Ibom State, Nigeria

Email: imehokop@aksu.edu.ng

Orcid id: [0000-0002-1487-6711](https://orcid.org/0000-0002-1487-6711)

1.0 Introduction

Piper guineense is an erect herbaceous climbing liana native to tropical Africa, *P. guineense* fruits are widely applied externally as a counter-irritant or in a stimulating ointment, internally as a stomachic and carminative. The leaves are useful in the treatment of wounds while the stems and twigs are for the treatment of coughs and bronchitis (Owolabi *et al.* 2013). Food and food materials can be assessed based on its chemical composition (Eko and Eddy, 2006). According to Imo *et al.* (2018), *Piper guineense* seeds have higher percentage of dry matter (94.03 ± 0.21), crude lipid (4.06 ± 0.12) and carbohydrates (65.46 ± 0.85) than the leaves while the leaves have higher percentage moisture (6.11 ± 0.01), protein (15.17 ± 0.39), crude fibre (20.99 ± 0.16) and ash (11.98 ± 0.03) than the seeds. Several extract of *Piper guineense* have been found to exhibit pharmaceutical and medicinal values. For example, methanol extract of *Piper guineense* was found to offered protection against infection that is comparable to that of Livolin forte with better efficacy when pre-treated with 400 mg/kg for 14 days prior to CCl₄-exposure (Oyinloye *et al.*, 2017). Ekudayo *et al.* (1988) identified elemicin as the major essential oil constituent of the plant and stated that the plant has significant medicinal applications. Olonisakin *et al.* (2006) identified (1s)-(-)- α -pinene (43.9%), D-Limonene (7.7%), caryophyllene (6.9%), car-2-ene (5.4%) and 1,6,10-dodecatrien-3-ol, 3, 7, 11-trimethyl (2.9%) and found that they did not display any antimicrobial activity against *Escherichia coli*, *Serratia*, *Salmonella typhi*, *Klebsiella sp.*, *Citrobacter* and *Pseudomonas aeruginosa* due to the solvent he used. Chinwendu *et al.* (2016) identified alkaloids (0.86%), HCN (8.87%), saponins

(1.87%) and phenols (0.66%) in ethanol extract of *Piper guineense* leaves while Ebenso *et al.* (2008) also identified the presence of alkaloids, tannins, saponins, flavonoids, hydrogen cyanides and phenols in ethanol extract of the leaves. They observed that, *Piper guineense* (Uziza leave) contains some considerable amount of anti-nutrients which have medicinal benefits.

GCMS analysis of plant extract has been found to be one of the most powerful tool that is useful for identifying chemical constituents of plants (Eddy *et al.*, 2011a,b; Ikpeazu *et al.*, 2020). Ojinnaka *et al.* (2016) identified 22 peaks from the GCMS spectrum of ethanol extract of *Piper guineense* leaves and found that the spectrum was dominated by acids and hydrocarbon while alcohol and ester were the least constituents. Recent study conducted by Usaman *et al.* (2020) indicated that the component extracted from plant parts depends on the type of solvent. However, most studies on *Piper guineense* leaves are done with aqueous and ethanol extract. Therefore, the present study is aimed at identifying the chemical constituents of acetone extract of *Piper guineense* leaves using GCMS analysis.

2.0 Materials and Methods

Samples of *Piper guineense* leaves were purchase from Ikot Ekpene main market and transported to the Chemistry laboratory of the Michael Okpara University of Agriculture, Umudike. They were thoroughly washed with distilled water and allowed to dry. The leaves were sun dried for a week until the moisture content was reduced to minimum. The dried leaves were grounded to a powder form and soaked in acetone solution. The solvent was recovered using cold extractor, leaving behind, acetone extract of *Piper guineense* leaves.

The produced extract was used for GCMS analysis using spectroscopically pure acetone solvent. The GCMS-QP2010 PLUS Shimadzu (made in Japan) instrument was used for the analysis. The analytical steps taken were plugger speed (high), syringe injection speed (high), viscosity/compression time (0.2 second), injection mode (normal), pumping time (5), injection port dwell time (0.3 second), terminated air cap (No), plugger washing speed (high), washing volume (8µl), syringe suction position (0), syringe injection position (0) and used three solvent vial (3). The operational setting of the GCMS instrument were column oven temperature (60°C), injection temperature (200°C), injection mode (split), flow control mode (linear velocity), pressure (100.2

kPa), total flow (6.2 ml/minute), linear velocity (46.3 cm/sec), purge flow (3.0ml/min) and split ratio (1.0). The high-pressure injection, carrier gas server and splitter hold functions were switch off. The initial rate of oven temperature program was 5 °C/min and was gradually increased to 140°C after which the temperature was increased to 280 °C at a rate of 10 °C/minute. Some heat unit and detector functions were checked in order to ensure consistency. These included column oven, SPL2, MS, SPL2 carrier, SPL2 purge and were ensured to be on. However, the APC setting was turned off.

Other setting functions of the machine were ion source temperature (200 °C), interface temperature (250 °C), solvent cut time ((2.50 minutes), detector gain mode (relative), detector gain (0.00kV), threshold (1000). The analytical start time was 3 minutes and the machine run for 45 minutes using ACQ scan mode at a scan speed of 769. However, mass/charge started at 50 and ended with 400 units.

Gas chromatogram and mass spectrum were automatically plotted and suggested chemical structures were obtained using the National Science Technology library installed in the machine. Percentage concentrations of each identified component was calculated using area normalization.

3.0 Results and Discussion

Fig. 1 presents the GCMS of acetone extract of *Piper guineense* while chemical names, retention time, molar mass and percentage concentrations of compounds deduced from each peak in the spectrum are recorded in Table 1. Fig. 2 shows the mass spectrum of the compounds.

The chemical constituents observed from the GCMS spectrum of the plant leaves were 3,6-dimethyloxazolo(5,4-c)pyridazin-4-amine (31.80 %), 3-(1-methylethyl)-cyclohexene (20.99 %), 4-methoxy-N-(4-nitrobenzyl)-benzamide (12.82 %), alpha bisabolene (7.33%), 1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methylene-1-(1-methylethyl)-naphthalene (4.42 %), 3,7-diacetamidophenoxathin (4.10 %), 1,3,3-trimethyl tricyclo{2,2,1}heptane (2.98%), 3H-indazol-3-one (2.11%), 1H-indene,octahydro-1,7a-dimethyl-4-(1-methylethenyl)-1,4-methano-1H-indene (1.98%), piperidine (1.97%),2,4-disopropenyl-1-methyl-1-vinyl (1.70%), n-hexadecanoic acid (1.68%), eudesma-4[14],11-diene (1.27%), tricosenoic acid (0.77%), alpha cubebene (Hctclopenta[1,3]cyclopropal[1,2]benzene (0.71%), 2,6,6-trimethyl-3-(phenylthio)cyclohept-4-enol



(0.69%), bicyclo[7,2,0]undec-4-ene,4,11,11-trimethyl-8-methylene (caryophyllene (0.54%), 1H-cycloprop[azulen-4-ol,decahydro-1,1,4,7-tetramethyl-globulol (0.47%) and 2,6-dimethoxytoluene (0.30%).










Fig. 1: GCMS of acetone extract of *Piper guineense*




Table 1: Characteristics of compounds in acetone extract of *Piper guineense* leaves

Line No	Name	Mass peak	Retention time (minute)	Molar mass (g/mol)	Concentration (%)
1	2,6-dimethoxytoluene	32	15.333	152	0.30
2	alpha cubebene (H-ctclopenta[1,3]cyclopropal[1,2]benzene	40	17.517	204	0.71
3	2,4-disopropenyl-1-methyl-1-vinyl	62	19.308	204	1.70
4	1,3,3-trimethyl tricyclo{2,2,1}heptane (cyclofenchene)	45	20.033	136	2.98
5	bicyclo[7,2,0]undec-4-ene,4,11,11-trimethyl-8-methylene (caryophyllene)	65	20.108	204	0.54
6	Cis alpha bisabolene	82	21.133	204	7.33
7	eudesma-4[14],11-diene	69	21.308	204	1.27
8	1,2,3,4,4a,5,6,8a-octahydro-7-methyl-4-methylene-1-(1-methylethyl)-naphthalene (gamma murolene)	73	22.125	204	4.42
9	1,2,3,5,6,7,8,8a-octahydro-1,4dimethyl-7-(1-methylethenyl)-azulene	74	22.65	204	1.37
10	3-(1-methylethyl)-cyclohexene	114	25.658	124	20.99
11	1H-cycloprop(e) azulene-4-ol,decahydro-1,1,4,7-tetramethyl-globulol	72	25.775	222	0.47
12	1H-indene,octahydro-1,7a-dimethyl-4-(1-methylethenyl)-1,4-methano-1H-indene	80	27.375	204	1.98
13	2,6,6-trimethyl-3-(phenylthio)cyclohept-4-enol	72	28.65	262	0.69
14	n-hexadecanoic acid	89	32.283	256	1.68
15	Tricosenoic acid	94	34.067	352	0.77




16	3H-indazol-3-one	61	36.075	134	2.11
17	4-methoxy-N-(4-nitrobenzyl)-benzamide	139	37.95	286	12.82
18	3,6-dimethyloxazolo(5,4-c)pyridazin-4-amine	153	39.667	164	31.80
19	piperidine	130	40.467	285	1.97
20	3,7-diacetamidophenoxathin	172	43.75	314	4.10



**2,6-dimethoxytoluene****Alpha cubebene****2,4-disopropenyl-1-methyl-1-vinyl****1,3,3-trimethyl tricyclo{2,2,1}heptane (cyclofenchene)****caryophyllene****Cis-alpha bisabolene**


**Eudema-4[14]-11-diene****Gamma murulene****1,2,3,5,6,7,8,8a-octahydro-1,4dimethyl-7-(1-methylethenyl)-azulene**

CompName:Cyclohexene, 3-(1-methylethyl)- \$\$ 3-Isopropyl-1-cyclohexene # \$\$


**3-(1-methylethyl)-cyclohexene**

CompName:(-)-Globulol \$\$ 1H-Cycloprop[e]azulen-4-ol, decahydro-1,1,4,7-tetramethyl-, [1aR-(1a.alpha.,4.alpha.,4a.alpha.,7a.alpha.,7a.beta.,7b.alpha.)]- \$\$


**1H-cycloprop(e) azulene-4-ol,decahydro-1,1,4,7-tetramethyl-globulol****1H-indene,octahydro-1,7a-dimethyl-4-(1-methylethenyl)-1,4-methano-1H-indene**




2,6,6-trimethyl-3-(phenylthio)cyclohept-4-enol




n-hexadecanoic acid



Tricosenoic acid



3H-indazol-3-one



4-methoxy-N-(4-nitrobenzyl)-benzamide






Fig. 2: Mass spectrum and chemical structures of compounds in acetone extract of *Piper guneense*

The mass spectrum of 3,6-dimethyloxazolo(5,4-c)pyridazin-4-amine was shown in Fig. 2. The compound is also called octocrinsghpfl-uhfffaoy-s-N. Little is known of its bioactivity or other usefulness. However, the 5,4 derivatives of the compound have been implicated in the treatment of cancer tumor. Caryophyllene identified in line 5 (of the GCMS spectrum) contains several biological activities that are attributed to betacaryophyllene, such as anti-inflammatory, antibiotic, antioxidant, anticarcinogenic and local anaesthetic. Piperidine (peak 19) has been confirmed to be active as antibacterial, analgesic and also exhibited anti-inflammatory activity (Mohammed *et al.*, 2016). Alpha and beta cubebene were also reported in essential oils of *Annona salzmannii* and *A. pickelii* (Annonaceae) by Coataa *et al.* (2011) and were found to exhibit strong anti-bacterial activity. Boligon *et al.* (2012) also attributed antimicrobial activity of *Scutia buxifolia* Reissek

leaves to the presence of cubebene. Haznedaroglu *et al.* (2001) found that the essential oil of *Salvia tomentosa* contains 1,8-cineol (17%), β -caryophyllene (11%), cyclofenchene (10%) and δ -cadinene (6%). They attributed the antimicrobial activity of the essential oil to these constituents, which also inhibited the growth of tested Gram-positive and Gram-negative bacteria except for *Pseudomonas aeruginosa*. Alpha-cubebene, camphene, geraniol, limonene, myrcene, palmitic acid and sabinene were found to exhibit antioxidant (DPPH assay), anti-inflammatory (5-lipoxygenase assay), antimicrobial (disk diffusion) and anti-mosquito properties (insecticidal, larvicidal and repellency assays) by Naidoo *et al.* (2009). α -farnesene and bisabolene are known flavour ingredients and their catalytic hydrogenation gives the hydrocarbons, farnesane and bisabolane, respectively. These saturated derivatives are prospective industrial products as they have been



singled out among the most promising biofuel candidates (Clarke, 2008). Sun *et al.* (2005) isolated four eudema and found that the compounds showed glucose consumption activity with an IC value of 10.7 microg/mL in a C2C12 muscle cell assay. The MIC value of this compound (100 mg/kg) in a db/db mice model was found to be equivalent to that of metformin in vivo. Limberger *et al.* (2001) found gamma murolene (identified in line 8) in *Blepharocalyx salicifolius* and linked it to some biological activities. Silva *et al.* (2009) also reported that gamma murolene is active against *Bacillus subtilis* and *Candida tropicalis*, including clinical strains. Tan *et al.* (2008) investigated and obtained results which indicated that globulol (identified in line 11) is the main antimicrobial compound in the ethanol extract of *E. globulus* fruits. Hexadecanoic acid (palmitic acid), was identified in line 14 of the spectrum. It is a saturated long-chain fatty acid with a 16-carbon backbone which has been reported to have potential antioxidant, antitumor, anti-inflammatory, antibacterial and antifungal activities (Vasudevan *et al.*, 2012).

4.0 Conclusion

GCMS of acetone extract of *Piper guineense* leaves reveals that the plant leaves contain organic acids, hydrocarbon, alcohol, ester, ketone and other aromatic compounds. Twenty compounds were identified and some of the compounds found in the spectrum differs from those reported by others. Most of the the identified compounds have industrial application and exhibit significant biological activities.

5.0 References

Boligon AA, Schwanz TG, de Brum TF, Frohlich JK, Nunes L, et al. (2012) Chemical Composition, anti-oxidant and anti-microbial activities of the eessential oil of *Scutia buxifolia reissek* leaves. *Pharmaceutical Analytica Acta*, 3, 199, doi:10.4172/2153-2435.1000199

Chinwendu, S., Ejike, E. N., Ejike, B. U., Oti, W. I. & Nwachukwu, I. (2016). Phytochemical properties of uziza Leave (*Piper guineense*). *European Journal of Pure and Applied Chemistry*, 3, 2, pp. 12-15.

Clarke, S. (2008). Essential Chemistry for aromaticity, 2nd Edition, Elsevier Ltd. Churchill, Livinstone, <https://doi.org/10.1016/B978-0-443-10403-9.X0001-5>

Coata, E. V., Dutraa, L. M., de Jesusa, H. C. R., Nogueiraa, P. C. D., Moraesa, V. R. D., Salvadorb, M. J., avalcantic, S. C. D., dos Santosd, R.

L. & Pratae, A. P. D. (2011). Chemical composition and antioxidant, antimicrobial, and larvicidal activities of the essential oils of *Annona salzmannii* and *A. pickelii* (Annonaceae). *Natural Product Communication*, 6, 6, pp. 907 – 912.

Ebenso, E. E., Eddy, N. O. & Odiongenyi, A. O. (2008). Corrosion inhibitive properties and adsorption behaviour of ethanol extract of *Piper guineensis* as a green corrosion inhibitor for mild steel in H₂SO₄. *African Journal of Pure and Applied Chemistry*, 4, 11, pp. 107-115.

Eddy, N. O., Awe, F. E., Siaka, A., Magaji, L. & Ebenso, E. E. (2011b). Chemical information from GC-MS studies of ethanol extract of *Andrographis paniculata* and their corrosion inhibition potentials on mild steel in HCl solution. *International Journal of Electrochemical Sciences*, 6, pp. 4316-4328.

Ekop, A. S. & Eddy, N. O. (2006). Comparative studies of the lipid characteristics and industrial potential of *Coula edulis* (African walnut and *Terminalia catappa* (indian almond) seeds. *Global Journal of Pure and Applied Sciences*, 12, 1, pp. 65-67.

Eddy, N. O., Ameh, P., Gimba, C. E. & Ebenso, E. E. (2011a). GCMS studies on *Anogessus leocarpus* (AL) gum and their corrosion inhibition potentials for mild steel in 0.1 M HCl. *International Journal of Electrochemical Sciences*, 6, pp. 5815-5829.

Ekundayo, O., Laakso, I., Adegbola, R.M., Oguntimein, B., Sofowora, A. & Hiltunen, R. (1988). *Essential oil constituents of Ashanti pepper (Piper guineense) fruits (berries)*. *Journal of Agriculture and Food Chemistry*, 36, 5, pp. 880-882, <https://doi.org/10.1021/jf00083a001>

Haznedarogku, M. Z., Karabay, N. U. & Zeybek, U. (2001). Antibacterial activity of *Salvia tomentosa* essential oil. *Fitoterapia*, 72, 7, pp. 829-831

Ikpeazu, O. V., Otuokere, I. E. & Igwe, K. K. (2020). Gas Chromatography–Mass Spectrometric Analysis of Bioactive Compounds Present in Ethanol Extract of *Combretum hispidum* (Laws) (Combretaceae) Root. *Communication in Physical Sciences*, 5,3, pp. 325-337.

Imo, C., Yakubu, O. E., Imo, N. G., Udegbumam, I. S., Tatah, S. V. & Onukwugha, O. J. (2018). Proximate, mineral and phytochemical composition of *Piper guineense* seeds and leaves. *Journal of Biological Sciences*, 18, pp. 329-337.



- Limberger, M.E.G. Sobral, J.A.S. Zuanazzi, P.R.H. Moreno, E.E.S. Schapoval & A.T. Henriques (2001) Biological Activities and Essential Oil Composition of Leaves of *Blepharocalyx salicifolius*, *Pharmaceutical Biology*, 39:4, 308-311, DOI: 10.1076/phbi.39.4.308.5915
- Mohammed, G. J., Omran, A. M., & Hussein, H. M. (2016). Antibacterial and Phytochemical Analysis of *Piper nigrum* using Gas Chromatography – Mass Spectrum and Fourier-Transform Infrared Spectroscopy. *International Journal of Pharmacognosy and Phytochemical Research*, 8, 6, pp. 977-996
- Naidoo, N., Thangaraj, K., Odhav, B., & Baijnath, H. (2009). Chemical composition and biological activity of the essential oil from *cymbopogon nardus* (L.) rendle. *African Journal of Traditional, Complementary and Alternative Medicines*, 6, 395. Retrieved from <https://journals.athmsi.org/index.php/ajtcam/article/view/779>
- Ojinnaka, M. C, Ubbor, S. C., Okudu, H. O. & Uga, U. (2016). Volatile compound analysis of the leaves and seeds of *Piper guineense* using gas chromatography-mass spectrometry (GC-MS). *African Journal of Food Science*, 10, 11, pp. 327-332
- Olonisakin, A., Oladimeji, M. O. & Lajide, L.(2006). Chemical Composition and Antibacterial Activity of Steam Distilled Oil of Ashanti Pepper (*Piper guineense*) Fruits (Berries). *Electronic Journal of Environmental, Agricultural and Food Chemistry*, 5, 5, pp. 1531-1535.
- Owolabi, M. S., Lawal, O. A., Ogunwande, I. A., Hauser, R. M. & Setzer, W. N. (2013). Aroma chemical composition of *Piper guineense* Schumacher & Thonn. From Lagos, Nigeria: a new chemotype. *American Journal of Essential oil and Natural Products*, 1,1, pp. 7-40.
- Oyinloye, B. E., Osunsanmi, F. O., Ajiboye, B. O., Ojo, O. A., & Kappo, A. P. (2017). Modulatory Effect of Methanol Extract of *Piper guineense* in CCl₄-Induced Hepatotoxicity in Male Rats. *International Journal of Environmental Research and Public Health*, 14, 9, pp. 955, <https://doi.org/10.3390/ijerph14090955>
- Silva, D. R., Endo, E., Dias Filho, B. P., Aparicio, D., Cortez, G., Nakaruma, C. V., Svidzinski, T. I. A., Souza, A., Young, M. C. M., Tania, U. & Cortez, D. A. G. (2009). Chemical Composition and Antimicrobial Properties of *Piper ovatum* Vahl. *Plant Medica*, 70, 9, pp. 1171-1182.
- Sun, Z., Chen, B., Zhang, B. & Hu, C. (2005). Four new eudesmanes from *Caragana intermedia* and their biological activities. *Journal of Natural Products*, 67, 12, pp. 1975-9
- Tan, M., Zhou, L., Huang, Y., Wang, Y., Hao, X. & Wang, I. (2008). Antimicrobial activity of globulol isolated from the fruits of *Eucalyptus globulus* Labill. *Natural Product Research*, 22, 7, pp. 569-575.
- Usman, A., Mohammed, Y., Muhammed, H. O. & Zakari, A. H. (2020). Phytochemical Screening and Antioxidant Activity of *Balanites Aegyptiaca* Root Bark Extracts: Influence of solvent. *Communication in Physical Sciences*, 5, 2, pp. 152-164.
- Vasudevan, A., Vijayan, D., Mandal, P., Karthe, P., Sadasivan, C. & Haridas, M. (2012). Anti-Inflammatory Property of n-Hexadecanoic Acid: Structural Evidence and Kinetic Assessment. *Chemical Biology and Drug Design*, 80, 3, pp. 434-439.

